

**PECTIN**  
**IDENTIFICATION**  
PRODUCT INSTRUCTIONS

**SKU: 700004325**  
**K-PECID**

11/24

(500 Assays per Kit)



© 2024, Neogen Corporation; © 2024, Megazyme. All rights reserved.

Neogen is a registered trademark of Neogen Corporation. Megazyme is a registered trademark of Megazyme Ltd.

## **INTRODUCTION:**

Pectins consist of the partial methyl esters of polygalacturonic acid and their sodium, potassium, calcium and ammonium salts, obtained by extraction in an aqueous medium of appropriate edible plant material, usually citrus fruits or apples. Pectin is recovered from solution by precipitation with an appropriate organic solvent such as ethanol, methanol or isopropanol. In some pectins, a portion of the methyl esters may have been converted to primary amide groups by treatment with ammonia under alkaline conditions. Older methods for the identification of pectin involve solvent precipitation. These methods are non-specific. The method described here measures the increase in absorption at 235 nm on cleavage of pectate by a highly purified and specific enzyme, pectate lyase.<sup>1</sup> The enzyme employed since 2011 is still derived from *Aspergillus niger* but it is different from that employed up until this date (see Figure 2, page 6).

## **PRINCIPLE:**

Pectin is dissolved in deionised water and adjusted to pH 12.0 to catalyse demethylation with the production of polygalacturonic acid regions in the polymer, i.e. conversion of pectin to pectate. The pectate is incubated with pectate lyase which cleaves the polygalacturonic acid, releasing unsaturated oligosaccharides which absorb strongly at 235 nm. The JECFA method<sup>2</sup> for the identification of pectins is very similar to the current method, however, the pectate lyase supplied with this kit acts better at pH 8.0.

## **SAFETY:**

The general safety measures that apply to all chemical substances should be adhered to.

For more information regarding the safe usage and handling of this product please refer to the associated SDS that is available from the Megazyme® website.

## **KITS:**

Kits suitable for performing 500 assays are available from Megazyme and consist of:

1. Concentrated Pectate Lyase enzyme
2. Low ester pectin extracted from citrus peel
3. High ester pectin extracted from citrus peel
4. Partly amidated low ester pectin from citrus peel
5. Pectin from sugar beet pulp
6. Lota carrageenan

## **SPECIFICITY:**

The assay is specific for polygalacturonic acid.

## **ENCLOSED ENZYME:**

**Pectate Lyase** (2.7 mL, 1400 U/mL at pH 8.0; Tris buffer)  
(500 U/mL at pH 10.8; CAPS buffer).

Store below -10°C. See individual label for expiry date.

**For use in the assay** of pectin, dilute 0.5 mL of enzyme to 50 mL in Tris-HCl buffer (pH 8.0) (i.e. a 100-fold dilution). Once dissolved, the enzyme is stable for 2 years below -10°C and to multiple freezing and thawing cycles.

## **ENCLOSED STANDARDS:**

1. Low ester pectin extracted from citrus peel
2. High ester pectin extracted from citrus peel
3. Partly amidated low ester pectin from citrus peel
4. Pectin from sugar beet pulp
5. Lota carrageenan

## **BUFFERS AND REAGENTS:**

1. **50 mM Tris/HCl buffer plus 1 mM CaCl<sub>2</sub>**  
Dissolve 6.055 g of Tris buffer (Megazyme cat. no. **B-TRIS500**), and 0.147 g of calcium chloride dihydrate in 900 mL of deionised water. Adjust pH to 8.0 with 1 M HCl. Adjust volume to 1 L.
2. **0.5 M NaOH**  
Dissolve 20 g of NaOH in 1 L of deionised water.
3. **0.5 M HCl**  
Add 50 mL of concentrated HCl (10 M) to 950 mL of deionised water.
4. **1 M HCl**  
Add 100 mL of concentrated HCl (10 M) to 900 mL deionised water.
5. **2-Propanol (100%)**

### SAMPLE PREPARATION:

1. Moisten 50 mg (0.05 g) of the sample with 2 drops of 2-propanol.
2. Add 50.0 mL of deionised water and stir gently on a magnetic stirrer for 20-30 min (until the pectin dissolves).
3. Adjust the pH to 12 by careful addition of 0.5 M NaOH, and leave the solution for exactly 15 min at room temperature.
4. Lower the pH to 8.0 by dropwise addition of 0.5 M HCl.
5. Adjust the volume to 100.0 mL with deionised water.

Add the following to quartz cuvettes, mix the contents well and measure the absorbance values at 235 nm after 30 min.

	Tris-HCl buff. (pH 8.0)	Sample	Deionised water	Diluted enzyme
enzyme blank	0.5 mL	1.0 mL	1.0 mL	-
sample blank	0.5 mL	-	1.5 mL	0.5 mL
reaction	0.5 mL	1.0 mL	0.5 mL	0.5 mL

### RESULTS:

The increase in absorbance for a given sample on incubation with pectate lyase is measured as follows:

**Blank Absorbance** = Enzyme Blank + Sample Blank  
(measured after 30 min)

**$\Delta$  Absorbance** = Reaction Absorbance – Blank Absorbance

From the increase in absorbance ( $\Delta A$ ) the amount of unsaturated product produced can be calculated as:

**Unsaturated product** =  $\Delta A/L \quad \times \quad 1/\epsilon$

**where:**

$\Delta A$  = Reaction Absorbance (after 30 min) – Blank Absorbance.


L = path-length of the reaction cuvette (= 1 cm)

$\epsilon$  = the molar extinction coefficient of the reaction product ( $4600 \text{ M}^{-1} \text{ cm}^{-1}$ )

This assay can be performed in a standard UV spectrophotometer simply by measuring the absorbance values of the reaction solution and the two blank solutions after the 30 min incubation.

However, the reaction can also be followed in a recording UV spectrophotometer set-up for kinetic assays. The results for such a study are shown in Figure 1.

**Figure 1.** Increase in absorbance at 235 nm on incubation of pectic polysaccharides (or carrageenan) with pectate lyase.



**Incubation conditions:**

1.0 mL of sample (0.5 mg/mL) + 0.5 mL deionised water + 0.5 mL Tris-HCl buffer (pH 8.0, containing 1 mM CaCl<sub>2</sub>).

**Add:** 0.5 mL of diluted pectate lyase (14 U/mL).

**Mix** immediately and follow the absorbance increase at 235 nm in a recording spectrophotometer thermostated at 40°C.

**REFERENCES:**

1. Hansen, K. M., Thuesen, A. B. & Soderberg, J. R. (2001). Enzyme assay for identification of pectin and pectin derivatives, based on recombinant pectate lyase. *J. AOAC International*, **84**, 1851-1854.
2. Compendium of Food Additive Specifications, FAO Food and Nutrition paper 52, Add. 9, Joint FAO/WHO Expert Committee on Food Additives (JECFA) 57th session, Rome, Italy, 5-14th June, 2001.

Polysaccharide type	Absorbance values				Unsaturated product x 10 <sup>-4</sup>
	Enzyme Blank	Sample Blank	Sample Reaction	ΔA	
Amidated low ester pectin	0.005	0.049	0.910	0.856	1.86
Sugar beet pectin	0.005	0.049	0.593	0.539	1.17
Carrageenan	0.003	0.047	0.052	0.002	0.00
High ester pectin	0.004	0.047	1.117	1.066	2.32
Low ester pectin	0.001	0.050	1.263	1.212	2.63

**Table 1.** Determination of the content of unsaturated oligosaccharides in pectic and non-pectic polysaccharides



Contact us for more information: [neogen.com/contact](https://neogen.com/contact)

---

**Without guarantee**

The information contained in this assay protocol is, to the best of our knowledge, true and accurate, but since the conditions of use are beyond our control, no warranty is given or is implied in respect of any recommendation or suggestions which may be made or that any use will not infringe any patents. It is the user's responsibility to perform in-house matrix validation work prior to routine use.

© 2024, Neogen Corporation; © 2024, Megazyme. All rights reserved.

Neogen is a registered trademark of Neogen Corporation. Megazyme is a registered trademark of Megazyme Ltd.